

Full Length Research Paper

Wild medicinal plants and their usage in traditional human therapy (Southern Bosnia and Herzegovina, W. Balkan)

Sulejman Redzic

Department of Botany, Faculty of Science, University of Sarajevo, Bosnia and Herzegovina.
E-mail: redzic0102@yahoo.com. Tel/Fax: +387 33 649 196.

Accepted 9 April, 2010

Both Mediterranean and sub-Mediterranean regions of Bosnia and Herzegovina are known for very prominent traditional usage of wild medicinal and aromatic plants for the treatment of various diseases. Ethno botanical interview method was used to collect information from the sample population comprised of 55 adults, of different religious background (Catholics, Muslims, and Orthodox), with an average age of 63. This resulted in the collection of 96 wild plants from 46 different plant families. These plants are used in preparation of up to 200 different ethno pharmaceuticals and used for 430 different treatments, mainly of chronic diseases: respiratory system (63 species), stomach and intestinal system (55 species), liver and gall bladder (60 species), urinary system (33 species), genital system (42 species), nervous system (30 species), cardiovascular system (27 species), skin conditions (56 species). 60 species are being used for the treatment of some other health disorders. In ecological sense, 40% of these species are constituents of meadow and rock debris communities, 34% of termophilous forest and shrub communities, while 20% of species are constituents of abandoned places.

Key words: Biodiversity, botanical medicine, ethno therapy, ethno biology, ethno pharmacology, herbal anthropology, pharmaceutical ethno botany, preparations of traditional medicines.

INTRODUCTION

Traditional herbal medicine has played an important role in the life of the Bosnian and Herzegovinian population. This is especially true for Herzegovinian population (Redzic, 2007), whose geographical area is generally very rich in medicinal plants (Redzic, 2009). Medicinal plants of this area have been used in treatment of various diseases since ancient times (Gluck, 1892; Sadikovic, 1928). Etnobotanical research in Herzegovina is an important link to similar researches in the other parts of the world, including both, third world countries (Girach et al., 1998; Lev and Amar, 2002; Fernandez et al., 2003; Giday et al., 2003; Hattarai et al., 2006) and the developed ones (Kenny, 2002; Applequist, 2004; Stobart, 2005; Pieroni et al., 2005a). Men have always used natural resources of healing substances in a search for a cure for a many pathological human condition. Effort to heal the sicknesses by means of traditional phyto-therapy has been made in all parts of the world (Heinrich, 2003;

2005; Abella et al., 2000; Bodeker et al., 2005). At present, ethno botanical and ethno pharmacological experiences of certain nation are used in the treatment of wide range of diseases (Matavelle and Habib, 2000; Sheng-Ji, 2001; Vivienne et al., 2005; Muthu et al., 2006; Uniyal et al., 2006) including treatments for the difficult to cure diseases such as cancer, AIDS, Alzheimer's disease, alcoholism, etc. (Perry et al., 1999; Carai et al., 2000; Bedoya et al., 2001; Donaldson and Rex, 2004; Bailly et al., 2005; Heinrich and Bremner, 2006; Sajem and Gosai, 2006). Today, a major breakthrough in a fight against stress is reached with the phytotherapy that is based on traditional ethno botanical knowledge gained through experiences of almost all nations of the world (van-Wyk, 2002; Nigro et al., 2004; Norscia and Borgognini-Tarli, 2006).

Each geographical area is characterized by its unique biodiversity, people, culture, specific history and its living

needs. This results in many different forms of interaction between a man and environment, through the nourishment needs (Redzic, 2006a, b), and basic medicinal healing treatments (Redzic et al., 1991; 1998).

Isolation of some geographical regions over longer periods of time, due to the relief dynamism or a presence of large rivers, has led to the development of trends in usage of wild plants in medicinal treatment and supplementary nourishment (Leporatti and Corradi, 2001; Leporatti and Ivancheva, 2003; Bonet and Vales, 2003; Leduc et al., 2006). An area of Mediterranean and sub Mediterranean part of Bosnia and Herzegovina (BiH) named Herzegovina, extremely dynamic in its biodiversity, relief, climate, anthropogenesis and other factors, developed unique culture and tradition with specific ethno botanical and ethno pharmacological way of practice.

Even though Herzegovina has been inhabited since the ancient times and is known to possess a long tradition in ethno botany, there are scarce records on respective experiences concerning medicinal or edible wild plants. This paper points out certain aspects of traditional medicinal plants practice in phytotherapy by human community in the Mediterranean and sub-Mediterranean region of BiH.

The main aim of this paper is to make an inventory and to record medicinal and aromatic plants practices in human therapy in Mediterranean and sub-Mediterranean region of BiH. This paper also includes data gathering on preparatory procedures for some herbal drugs.

Study area

The area of interest is located between N 43° - 44° and E 17° - 19° (Figure 1). It includes Mediterranean and sub Mediterranean zone belt, as well as one part of Southern Herzegovinian high Mediterranean region. Mediterranean belt includes 30 km of Adriatic coastline. Sub Mediterranean belt extends to north towards Mostar city including area around Ljubuski in the west, and Stolac in the east. Altitude of the investigated area spans from 0 - 1,800 m above the sea level with an average altitude of around 300 m.

Natural vegetation of this area comprises of remnants of woods and shrubs with evergreen oak *Quercus ilex*, deciduous oak forests with *Quercus pubescens*, *Carpinus orientalis*, *Fraxinus ornus* and *Quercus frainetto*. In the colder habitats, vegetation includes shrubs of *Ostrya carpinifolia* and *Fagus moesiaca*. Characteristic of sub Mediterranean region is an occurrence of shrubs composed of endemic Balkan species *Petteria ramentacea* (Redzic, 1999; 2000). Secondary vegetation types include shrubs, xeric grasslands and meadows. Significant number of medicinal plants quite commonly grows in the habitats of the abandoned places (around roads and arable land). Moving towards the high mountain region, wood communities including the

endemic pine *Pinus heldreichii* can be found. This area is characterized by an enormous heterogeneity and the high level of habitat's diversity, as well as biodiversity in general. It also possesses richness of endemic and relict plants. 60% of 450 endemic plants of BiH are found in the territory of Herzegovina (Redzic et al., 2003).

Population

The territory of Herzegovina, around Neretva River, was inhabited by humans during a Neolithic Era (Fukarek, 1954) and their settlements were mainly located around a river beds. Some evidences prove existence of a pill dwelling. All investigated sites in the area of interest have (spanning from Adriatic Sea to Mostar city) revealed remains of the Roman culture and civilization. Later in history, these attractive areas were visited by many individuals arriving from the North, South, West, and East. The culture and tradition of the region were heavily influenced by the Austro-Hungarian monarchy and the Ottoman Empire. The area has been inhabited by Catholics and Muslims and later by Orthodox too. Similar population structure is maintained nowadays. This is confirmed by an existence of many sacral places, that are known beyond borders of BiH - Medjugorje (Sveta Gospa, Catholic), Manastir Žitomislici (Orthodox), and Tekija on Buna river (Islamic). Since the early days of civilization, the population was orientated to agriculture- (growing vegetables and Mediterranean fruits), and at higher altitudes - cattle breeding, especially goats. Since ancients, vine and tobacco were also grown here (Begovic, 1960). Over the last twenty years tangerine, kiwi, tomato, sweet pepper, watermelon, peach, and apricot have been planted as well.

MATERIALS AND METHODS

The intensive ethno botanical research has been conducted continuously from a year 2000 to the year 2005. Significant amount of data was gathered during the long-termed ethno botanical and floristic investigations in this part of BiH, which was carried out by the author during the last 15 - 20 years. They include all important transects (Figure 1): from the Adriatic Sea coast to Hutovo blato (17 informants), surrounding of Stolac (9 informants), surrounding of Ljubuski, villages Vitina and Klobuk (12 informants), surrounding of Čapljina (6 informants) and surrounding of Mostar, Podveležje and Buna (11 informants).

Ethno botanical interview has been used as a basic method for data gathering. All informants have given their consent to participate in the research. The ethno botanical interview form contained: name and age of informants, area/village, time the interview took place, local name of medicinal herbs, herbal part being used, preparation procedure, purpose of usage, habitat type and estimation of conservation status.

The survey included interviews of over 55 informants. Average age of informants is 63 (ranging from 48 - 83). They had to be involved in collection and application of medicinal plants with the therapeutic purpose in any way. Informants were identified through the local community authorities. Local community's representatives provided data on persons which could have certain experience in




Figure 1. Investigated area.

human therapy. The author made an oral interview with each of the above mentioned persons, explaining the research's objective and making a preliminary testing of their knowledge in the field of traditional therapy. Only after they gave their consent, they were

interviewed by the author. Printed forms containing following information: common or the most frequently used plant name, plant's part being used, usage modes and the purpose of usage were distributed to all informants. Thereafter, the author collected indicated plants and confirmed their true identity with informants. In some cases, the informants were able to provide only dried plant's materials and the author himself had to conduct further botanical identification. Knowledge on medicinal herbs and traditional use thereof in therapy and ethnology, was transferred from generation to generation. In the Mediterranean area people use wild plants for healing purposes more often than in any other region. Some informants stated to practice traditional healing modes with herbs based on ethno botanical and ethno pharmacological experiences gained by earlier herbalists. Many of them are very famous and respected, even today and their work has also been acknowledged by pharmacological scientists too, such as St. Grga Martić and Sadik Sadiković.

The informants are the members of different ethnical groups. Approximately 55% of them are Catholics, 35% are Muslims and 10% are Orthodox. Gender ratio is: approx. 65% female and 35% of male. In many occasions, the author presented the collected herbal material the locals, and for the each specimen that was recognized as the one that is being used in the therapy, relevant data was collected (including photos and prepared plant material).

Communication with the locals went on smoothly due to the fact that the entire population of the investigated area uses the same local language. Moreover, all interviewed persons, with the few exceptions, readily agreed to participate in ethno botanical dialog. In the dialogs, the author expressed the deepest respect to each interviewed person, especially to the respective ethno botanical skills and knowledge. High level of co-operation was also achieved thanks to the fact that the author is well known to the locals for his numerous TV presentations about nature and medicinal herbs.

The entire plant material collected in the course of research is placed and stored in the Herbaria of medicinal plants of the Center of Ecology and Natural Resources at the Faculty of Science, University of Sarajevo (CEPRES HERB!) and in the Herbaria of the National Museum of BiH (SARA!). Each determined specimen is assigned with own herbarium number (voucher of specimen).

Herbal material was identified by the author. Some questionable taxa were determined by comparing them with the referring specimens from SARA and CEPRES HERB. The plant nomenclature complies with Flora Europea (Tutin et al., 1980). It is important to point out that in some cases several modes of usage and preparation procedures for certain herbs were recorded. As a consequence of this, spectrum of occurring species concerns only the most frequent ones, whereby less frequent species were named in Table 1. Forms of usage of medicinal herbs used for the treatments have been created in consultation with the experienced local herbalists taking into account human organ systems. Here, like in other areas, some species, or even genera are highly variable and complex for precise taxonomic determination, especially in the sense of a modern taxonomic investigation (EURO-MED Plant Base, 2006). Therefore, some species are aggregates that contain some micro-species whose origin are found within these aggregates. Example of these are *Achillea millefolium* agg., *Thymus serpyllum* agg., *Taraxacum officinale* agg., *Rosa canina* agg., *Rubus heteromorphus* agg.

RESULTS

During the course of investigations 96 vascular plants have been recorded belonging to the wild flora and playing a key role in the phytotherapy of Mediterranean and Sub-Mediterranean region of BiH. An overview of

Table 1. Wild medicinal flora of Mediterranean and Sub-Mediterranean region of Bosnia and Herzegovina use in ethno therapy.

Habitat/belonging to plant community*	Scientific name of plant, plant family and voucher herbarium specimens	Local name	English name	Part (s) used	Frequency of citation
Arr./T.-H.	<i>Achillea millefolium</i> L. Asteraceae (210101H)	Kkkkkkkkk Kunica , sporiš	Yarrow	Aerial part	26
Ph.	<i>Acrus calamus</i> L. Araceae (220504H)	Idjirot	Sweet flag	Rhizome	5
Ad.	<i>Adiantum capillus-veneris</i> L. Adiantaceae (310101H)	Vilina vlas	Maidenhair fern	Aerial part	2
T.-H.	<i>Althaea officinalis</i> L. Malvaceae (210601H)	Bijeli sljez	Marsh mallow	Leaf, Root	12
B.e.	<i>Anthyllis vulneraria</i> L. Fabaceae(210901H)	Ranjenik	Kidney vetch	Aerial part	4
Q.i.	<i>Arbutus unedo</i> L. Ericaceae (210580H)	Planika	Strawberry tree	Leaf, fruit	10
O.	<i>Arctium lappa</i> L. Asteraceae (210103H)	Čičak, repuh	Great burdock	Root, leaf, fruit	13
O.	<i>Artemisia absinthium</i> L. Asteraceae (210104H)	Pelin, pelim	Wormwood	Aerial part	21
O.	<i>Artemisia vulgaris</i> L. Asteraceae (210105H)	Komonjika	Mugwort	Young shoots	7
O.-C.o.	<i>Asparagus officinalis</i> L. Asparagaceae (220605H)	Viline metle	Asparagus	Young shoots	7
Q.p.	<i>Calamintha officinalis</i> Moench Lamiaceae (210503H)	Marulja	Calamint	Aerial part	5

Table 1. Contd.

Ch.	<i>Capsella bursa-pastoris</i> (L.) Med. <i>Brassicaceae</i> (210151H)	Rusomača	Shepherd's purse	Aerial part	11
B.e.	<i>Carlina acaulis</i> L. <i>Asteraceae</i> (210108H)	Kravljak, sikavac	Stem less carline thistle	Root	7
C-B.r.	<i>Centaurium maritimum</i> (L.) Fritsch <i>Gentianaceae</i> (210590H)	Kantarija, kitica	Centaury	Aerial part	10
Amp.	<i>Ceterach officinarum</i> DC. <i>Aspleniaceae</i> (310102H)	Zlatna paprat	Scale fern	Leaf	7
Ch.	<i>Chelidonium majus</i> L. <i>Papveraceae</i> (210190H)	Rosopas	Greater celandine	Aerial part	16
Sis.	<i>Cichorium intybus</i> L. <i>Asteraceae</i> (210110H)	Vodopija, ženetrğa	Chicory	Root	14
Sis.	<i>Cnicus benedictus</i> L. <i>Asteraceae</i> (210111H)	Blaženi čkalj	Blessed Thistle	Leaf	6
O.-C.o.	<i>Cotinus coggygria</i> Scop. <i>Anacardiaceae</i> (210220H)	Ruj	Smoke tree	Leaf	5
O.-C.o./ P.s.	<i>Crataegus monogyna</i> Jacq. <i>Rosaceae</i> (210403H)	Jednosjemeni glog	Hawthorn	Flower, leaf	31
Sis.	<i>Cynoglossum officinale</i> L. <i>Boraginaceae</i> (210240H)	Mišinac, mali gavez	Hound's tongue	Aerial part, leaf, root	7
Q.p.	<i>Dictamnus albus</i> L. <i>Rutaceae</i> (210471H)	Jasenak	Burning bush	Aerial part, root	2
P.m.	<i>Equisetum arvense</i> L. <i>Equisetaceae</i> (310201H)	Preslica	Field horsetail	Aerial part	7

Table 1. Contd.

B.e.	<i>Eryngium amethystinum</i> L. <i>Apiaceae</i> (210705H)	Kotrljan	Blue eryngo	Root	10
B.e.	<i>Filipendula vulgaris</i> Moench <i>Rosaceae</i> (210407H)	Kraljica polja	Dropwort	Aerial part	3
Sis.	<i>Foeniculum vulgare</i> Mill. <i>Apiaceae</i> (210706H)	Komorač	Fennel	Aerial part	17
P.s.	<i>Fragaria vesca</i> L. <i>Rosaceae</i> (210406H)	Jagoda šumska	Wild strawberry	Leaf, rhizome, fruit	15
O.-C.o.	<i>Fraxinus ornus</i> L. <i>Oleaceae</i> (210341H)	Crni jasen	Manna ash	Leaf	5
Ch.	<i>Fumaria officinalis</i> L. <i>Papaveraceae</i> (210200H)	Dimnjača	Fumitory	Aerial part	3
A.f.	<i>Geranium macrorrhizum</i> L. <i>Geraniaceae</i> (210350H)	Zdravac planinski	Big root geranium	Rhizome	3
T.-H..	<i>Glechoma hederacea</i> L. <i>Lamiaceae</i> (210504H)	Dobričica	Ground Ivy	Aerial part	6
T.-H.	<i>Glycyrrhiza glabra</i> L. <i>Fabaceae</i> (210903H)	Sladić	Liquorices	Root	7
Q.p.	<i>Hedera helix</i> L. <i>Araliaceae</i> (210360H)	Bršljan	Ivy	Leaf	7
S.-Ch.	<i>Helichrysum italicum</i> (Roth.) G.Don <i>Asteraceae</i> (210112H)	Smilje	Curry plant	Flower	10
S.-Ch.	<i>Herniaria hirsuta</i> L. <i>Caryophyllaceae</i> (210370H)	Sitnica, kilavica	Hairy rupture wort	Aerial part	9

Table 1. Contd.

B.e.	<i>Hieracium pilosella</i> L. Asteraceae (210113H)	Runjika, zečija loboda	Mouse-Ear hawkweed	Aerial part, root	7
B.e.	<i>Hypericum perforatum</i> L. Clusiaceae (210390H)	Bogorodična trava	St. John's Wort	Aerial part	27
S.-Ch.	<i>Hyssopus officinalis</i> L. Lamiaceae (210505H)	Miloduh, isop	Hyssop	Aerial part	12
Q.p./P.a.	<i>Juglans regia</i> L. Juglandaceae (210400H)	Orah	Walnut	Leaf, immature fruits	9
Jun.	<i>Juniperus oxycedrus</i> L. Cupressaceae (110302H)	Crvena kleka	Prickly Juniper	Fruit	5
S.-Ch/Ch	<i>Lavandula vera</i> DC. Lamiaceae (250531H)	Despik	Lavender	Aerial part	11
Q.p.	<i>Lilium cattaniae</i> (Vis.)Vis. Liliaceae (220503H)	Zlatan	Balkan Lily	Bulb	2
Ch.	<i>Malva neglecta</i> Wallr. Malvaceae (211231H)	Mali sljez	Dwarf Mallow	Aerial part	7
S.-Ch.	<i>Marrubium incanum</i> Dsr. Lamiaceae (210507H)	Matočina, očajnica	SIVI Horehound	Aerial part	7
Q.p.	<i>Melittis melissophyllum</i> L.. Lamiaceae (250528H)	Medunika	Bastard Balm	Aerial part	4
T.-H.	<i>Mentha pulegium</i> L. Lamiaceae (210511H)	Verem trava	Pennyroyal	Aerial part	15
S.-Ch./ Amph.	<i>Micromeria thymifolia</i> (Scop.) Fritsch. Lamiaceae (210512H)	Timjanoliki vrisić	Thyme Savory	Aerial part	3

Table 1. Contd.

O.	<i>Nepeta cataria</i> L. Lamiaceae (210513H)	Mačija metvica	Catmint	Aerial part	17
Ch	<i>Nigella damascena</i> L. Ranunculaceae (210205H)	Čurekot, mačkov brk	Love-In-A-Mist	Seed	2
B.e.	<i>Orchis morio</i> L. Orchidaceae (220502H)	Salep, kačun	Green-Winged Orchid	Tuber	17
O.-C.o.	<i>Origanum heracleoticum</i> L. Lamiaceae (250530H)	Mravinac	Oregano	Aerial part	10
Pal./ O.-C.o.	<i>Paliurus spina-christi</i> Mill. Rhamnaceae (210294H)	Drača	Christ's thorn	Fruit	2
B.e.	<i>Pimpinella saxifraga</i> L. Apiaceae (210708H)	Bedrenika	Burnet Saxifrage	Root	6
P.h.-n.	<i>Pinus heldreichii</i> H. Christ Pinaceae (110103H)	Munika		Resin	2
P.h.-n.	<i>Pinus nigra</i> J.F. Arnold Pinaceae (110105H)	Crni bor	Austrian pine	Resin	10
Q.i.	<i>Pistacia lentiscus</i> L. Anacardiaceae (210145H)	Tršlja	Mastic tree	Leaf, bark	3
B.e.	<i>Plantago media</i> L. Plantaginaceae (210461H)	Srednja bokvica	Hoary plantain	Leaf	10
Q.p.	<i>Polygonatum odoratum</i> (Mill.) Druce Convallariaceae (220604H)	Pokosnica	Solomon's seal	Rhizome	5
P.m.	<i>Polygonum aviculare</i> L. Polygonaceae (210321H)	Troskot	Knotweed	Aerial part	12

Table 1. Contd.

Amph.	<i>Polypodium vulgare</i> L. <i>Polypodiaceae</i> (310103H)	Slatka paprat	Polypody	Rhizome	4
T.-H.	<i>Potentilla anserine</i> L. <i>Rosaceae</i> (210416H)	Guščija trava	Polypody	Leaves	3
B.e.	<i>Prunella laciniata</i> (L.) L. <i>Lamiaceae</i> (250531H)	Bijela celinčica	Self-Heal	Aerial part	4
C.-B.r.	<i>Psoralea bituminosa</i> L. <i>Fabaceae</i> (210904H)	Djetelinjak	Silver leaf	Aerial part	2
O.-C.o.	<i>Punica granatum</i> L. <i>Punicaceae</i> (210300H)	Nar	Pomegranate	Bark	5
Q.p.	<i>Quercus pubescens</i> Willd. <i>Fagaceae</i> (210273H)	Hrast medunac	Downy Oak	Bark, fruit	7
O.-C.o.	<i>Rhamnus cathartica</i> L. <i>Rhamnaceae</i> (210290H)	Pasdrijen	Common buckthorn	Bark	6
P.s.	<i>Rosa canina</i> L. <i>Rosaceae</i> (210410H)	Šipurak	Dog rose	Fruit	12
Q.i.	<i>Rubia peregrina</i> L. <i>Rubiaceae</i> (210112H)	Broćac	Wild madder	Aerial part	3
O.-C.o. /P.s.	<i>Rubus heteromorphus</i> Ripart ex Genev. <i>Rosaceae</i> (210415H)	Dalm.kupina	Dalmatic blackberry	Leaf, fruit	9
O.-C.o.	<i>Ruscus aculeatus</i> L. <i>Liaceae</i> (220605H)	Veprina	Butcher's broom	Rhizome	3
S.-Ch.	<i>Ruta chalepensis</i> L. <i>Rutaceae</i> (210470H)	Ruta, sedef	Egyptian rue	Aerial part	9

Table 1. Contd.

P.a.	<i>Salix alba</i> L. Salicaceae (210332H)	Bijela vrba	White willow	Bark	6
B.e.	<i>Salvia bertoloni</i> Vis. Lamiaceae (210529H)	Kadulja livadska	Grassland sage	Leaf	8
S.-Ch.	<i>Salvia officinalis</i> L. Lamiaceae (210528x)	Kadulja	Sage	Leaf	31
O.	<i>Sambucus ebulus</i> L. Caprifoliaceae (210480H)	Apta, burjan	Dwarf Elder	Leaf, root, fruit	5
B.e.	<i>Sanguisorba minor</i> Scop. Rosaceae (210417H)	Dinjica	Salad Burnet	Fresh leaf	9
S.-Ch.	<i>Satureia montana</i> L. Lamiaceae (210516H)	Vrijesak bijeli	Winter Savory	Aerial part	12
S.-Ch.	<i>Satureia subspicata</i> Bartl. ex Vis. Lamiaceae (210517H)	Vrijesak crveni	Red Savory	Aerial part	12
S.-Ch.	<i>Sedum acre</i> L. Crassulaceae (210490H)	Žuti žednjak	Common Stonecrop	Aerial part	11
Amph.	<i>Sedum maximum</i> (L.) Suter Crassulaceae (210491H)	Debelo more	Orpine	Fresh leaf	15
S.-Ch.	<i>Sedum sexangulare</i> L. Crassulaceae (210492H)	Bradajušćak	Small Stonecrop	Fresh leaf	5
T.-H.	<i>Symphytum officinale</i> L. Boraginaceae (210243H)	Gavez	Comfrey	Root, leaf, fresh juice	21
Arr./O.	<i>Taraxacum officinale</i> Weber Asteraceae (210120H)	Maslačak, žučanik	Dandelion	Root, leaf, flower	31

Table 1. Contd.

A.f.	<i>Teucrium arduini</i> L. Lamiaceae (210519H)	Velika iva	Arduin's Germander	Aerial part	2
S.-Ch.	<i>Teucrium montanum</i> L. Lamiaceae (210521H)	Iva trava	Common Germander	Aerial part	21
C.-B.r.	<i>Teucrium polium</i> L. Lamiaceae (210522H)	Iva mediteranska	Mediterranean Germander	Aerial part	2
B.e.	<i>Thymus serpyllum</i> L. Lamiaceae (210524H)	Čubra, majkina dušica	Wild Thyme	Aerial part	32
S.-Ch.	<i>Thymus pulegioides</i> L. Lamiaceae (210525H)	Timijan, majkina dušica	Broad-Leaved Thyme	Aerial part	23
Q.p.	<i>Tilia cordata</i> Miller Tiliaceae (210500H)	Sitnolisna lipa	Small Leaved Lime	Flower	31
O.	<i>Tussilago farfara</i> L. Asteraceae (210122H)	Podbjel	Coltsfoot	Leaf, flower	23
O.	<i>Urtica dioica</i> L. Urticaceae (210510H)	Kopriva, žara	Stinging Nettle	Leaf, root, seeds	31
O.	<i>Verbascum thapsus</i> L. Scrophulariaceae (210256H)	Divizma, vučiji rep	Great Mullein	Leaf, Flower	5
O.-C.o.	<i>Viola alba</i> L. Violaceae (210552H)	Ljubičica bijela	White Violet	Flower, leaf	10
O.-C.o.	<i>Viola odorata</i> L. Violaceae (210551H)	Ljubica mirisna	Sweet Violet	Flower, leaf	12
Q.p.	<i>Viscum album</i> L. Violaceae (210560H)	Imela bijela	Mistletoe	Aerial part	10

Table 1. Contd.

S.p.	<i>Vitex agnus-castus L.</i> <i>Verbenaceae (210570H)</i>	Fratarski biber	Agnus Castus	Aerial part, seed	4
------	--	-----------------	--------------	-------------------	---

*Q.i. - *Quercetalia ilicis* Q.p.- *Quercetalia pubescentis* O.-C.o.- *Ostryo-Carpinetalia orientalis* P.s. -*Prunetalia spinosae* .Jun. - *Juniperetalia* Pal. - *Paliuretalia aculeatae* P.h.-n.- *Pinetalia heldreichii-nigrae* P.a. - *Populeetalia albae* S.p. - *Salicetalia purpureae* Arr. - *Arrhenatheretalia*, T.-H. *Trifolio-Hordetalia* B.e. - *Brometalia erecti* S.-Ch. - *Scorzonero-Chrysopogonetalia* C.-B.r. - *Cymbopogo-Brachypodietalia* Ph. - *Phragmitetalia* Ad. - *Adiantetalia* M.p. - *Moltkaetalia petraeae* (incl. Amph. - *Amphoricarpetalia*) D.j. - *Drypeetalia jacquiniana* (Incl. A.f. - *Arabidetalia flavescens*) Ch. - *Chenopodietalia* O. - *Onopordetalia* Sis. - *Sisymbrietalia* P.m. *Plantaginetalia majoris*

** E.r.–Edibility rating ***M.r.–Medicinal rating

Table 1. Contd.

Use in human therapy	Preparation and or/ Administration process
Wound, leucorrhea, chapped breast nipple, stomach, liver, hear palpitation, liver inflammation, gall disorder, purify of blood, strengthen of spleen and pancreas, hard diabetes	Fresh juice, infusion, oil
Fever, strengthen of nervous system, stomach, for better digestion, heart palpitation, cough, disagreeable felling, hepatitis	Infusion, powder
Plug, kidney, menstruation problem	Infusion
Cough, skin, tonic, voice improvement, fever, coldness, strengthen of liver, gall, inflammation of pancreas, hard hemorrhoids,	Maceration, light infusion
Wound, contusion, cough, skin disorders	Infusion
Kidney, urinary disorder, skin diseases, constipation	Decoction, fresh fruit
Skin, regeneration root of hair, with vine eliminate of tapeworm, melting of kidney stone, effective urination	Decoction, wine and "lozovaca" tincture
Diabetes, when to bite of fury dog, at stomach intoxication, venereal diseases, inflammation of facial nervous	
Asthma, Appetite, leucorrhea, heart disorders (arrhythmia), liver, gall, hepatitis	Infusion, tincture, powder
Leucorrhea, digestive disorders, water disease, psychotic disorders	Infusion, "lozovaca" Tincture
Urinary infection, prostate inflammation	Infusion
Sedative, tonic, cough, liver disorders	Infusion, tincture
Leucorrhea, Internal bleeding, skin disorders, Menstrual bleeding	Fresh juice, infusion
Catalyze of urination, strengthen of kidney, nervous system, at water disease, hepatitis, furuncle and skin diseases	Infusion, fresh juice, infusion in vinegar
Stomach disorders, recovery, genital disorder, leucorrhea	Infusion, tincture, infusion with honey
Urinary infections, skin diseases, cough	Infusion
Cough, asthma, constipation, Wart on skin, cancer of lung, hepatitis, against to uterus convulsions, melting of gall stone	Fresh juice, infusion, tincture
Water disease, anemia, inflammation of uterus, strengthen of liver and spleen	Infusion, tincture in lozovaca
Cancer of stomach, strengthen of liver and spleen	Infusion, tincture
Skin diseases, fever, hair regeneration	Infusion
Heart disease, sedative, relaxation	Infusion, tincture

Table 1. Contd.

Burn, bone fracture external, skin disorders	Maceration, decoction in milk
Sedative, stomach disorder, cough	Infusion
Furuncles, sin diseases, cough, melting of kidney stones, skin disorders, urinary infections, face pimples, inflammation of urinary bladder	Infusion
Water disease, hepatitis	Infusion
Rheumatism, circulation, fever, cough	Infusion, fresh juice
Cough, Stomach and respiratory disorders	Tincture, infusion, fresh juice
Recovery, diarrhea, strengthen of heart, melting of kidney stone and gallstone, ayes inflammation	Infusion, decoction
Constipation, fever, cough	Infusion, fresh fruit
Blood recovery, tonic	Infusion, fresh juice
Stomach disorder, menstruation disorders	Infusion, decoction
Respiratory system, hysteria, stone and sand kidney, hepatitis, infected stomach	Infusion, extract in oil olive
Asthma, Stomach disease, liver, skin disorders	Infusion, powder
Hepatitis, hard lung diseases, for better urination gall, skin disease	Infusion, extract in cooked wine
Liver, gall, cough, stomach and liver disorders	Infusion
Urinary system, kidney, prostate inflammation	Infusion
Urinary system, prostate gland inflammation	Infusion
Lung diseases, Stomach, skin disease, neurosis, kidney inflammation, venereal diseases, inflammation of spleen	Infusion, oil
Lung disease, tonsillitis, strengthen of potency	Infusion
Leucorrhea, liver disease, skin, strengthen, throat inflammation, strengthen of muscles, fore better breast , strengthen of male sex organs	Decoction, infusion
Asthma, skin disease, urinary infection, tuberculosis, rheumatism, gall stone melting, nervous disorders	Decoction, macerate, infusion, tincture in lozovaca
Asthma, uterus pains and inflammation, vaginal infections, rheumatism	Infusion, extract in olive oil
Wound, gall disorder, circulation	Extract in olive oil, infusion
Cough, skin disorder, catalyze of digestion, mouth diseases	Maceration, infusion
Anemia, heart disease, stomach disease	Infusion, tincture
Cough, stomach, sedative	Infusion
Vaginal infections, neural disease, headache	Infusion
Stomach and lung diseases, liver and gall disorders, against to hysteria, relaxation	Infusion, tincture
Hysteria, for barren woman, vaginal chronic infections	Infusion
Excretion of urine, prostate gland inflammation	Infusion, powder
Cough, sex power, constipation	Decoction
Cough, improvement potency, stomach, liver disorders	Infusion, tincture in lozovaca
Diarrhea, recovery, strengthen of stomach, skin disorders	Powder, decoction

Table 1. Contd.

Chronic vaginal infection, cough, stomach hardship purify of blood, strengthen of nervous system, regulate of menstruation, eliminate gas in stomach	Infusion, maceration
Contusion, wound, asthma, heart disorders	Resin, traditional balm
Wounds, rheumatism	Resin, traditional balm
Psoriasis, skin disorders	Decoction in wine
Hemorrhoids, bleeding from trachea, Cough, wound, liver disorders, uterus disorders, leucorrhea	Fresh juice, infusion
Bone fracture, skin diseases	Decoction, infusion
Vaginal infections, prostate gland, urinary infection, kidney inflammation, spleen, ulcer, hard cough, bleeding of lung	Infusion
Cough, skin disorder	Decoction, infusion
Asthma, diarrhea, liver, strengthen of heart	Decoction, infusion
Cough, liver, gall, hard skin disorder	Infusion
Constipation, skin disorder	Infusion
Diarrhea, stomach warm	Decoction, infusion
Vaginal chronic infections, recovery, skin disease, tonsillitis, fistulae and hemorrhoids	Decoction, maceration, powder
Chronic constipation, skin diseases, liver and Spleen disorders, hemorrhoids	Infusion, decoction
Diarrhea, roundworm, recovery, fever, cough	Infusion, maceration, decoction
Kidney disease, hepatitis	Decoction, infusion
Cough, diarrhea, against uterus bleeding, skin disorders, balance of body temperature	Decoction, juice
Hemorrhoids, blood vessels	Decoction, infusion
Heart disorder, against to hysteria and epilepsy	Fresh juice, infusion
Heart disorder, skin diseases, "swam fever" (malaria), corn at foot	Decoction, extract in wine
Skin disease, cough, liver and gall disorders	Infusion
Respiratory and skin disease, strengthen of nervous system, clean of liver, strengthen of kidneys, heard, inflammation of mouth mucous membrane, toothache, recovery and strengthen of organism	Infusion, extract in wine, tincture in lozovaca
Eliminate of toxic liquids by urine, clean of kidneys, water diseases, inflammation of urinary bladder,	Infusion, potion
Cough, Rheumatism, stomach disorder	
To bite of poisonous snakes, Diabetes, melting of kidney stone, skin disorders, diabetes	Fresh juice, infusion
Blood – leukemia, lung diseases, live, spleen, skin diseases, potency improvement	Infusion, powder, tincture in lozovaca
Blood disease, plug disease, potency, cough, liver and gall disorders, skin cleaning	Infusion, powder, tincture in lozovaca
Plug serious disease, skin wart, liver regeneration	Fresh juice
Liver, skin, wounds	Fresh leaves, mixture with honey
Chronic vaginal infections	Powder
Bone fracture, lung inflammation, asthma, bronchitis, kidney mucous elimination, baldness, hard fever	Decoction with milk, fresh juice
Anemia, Liver, diabetes, kidney stone melting, hemorrhoids, recovery, hepatitis	Infusion, decoction with milk

Table 1. Contd.

Stomach disease, liver and gall stone	Infusion, tincture in lozovaca
Liver, stomach disease, gall disorders	Infusion, tincture in lozovaca
Stomach disease, liver and gall disorders	Infusion, tincture in lozovaca
Neurosis, cough, stomach disease, irregular menstruation, strengthen of heart, stomach, anemia, hands tremor, paralysis of legs and hands	Infusion, powder, tincture in lozovaca, extract in olive oil
Neurosis, cough, stomach disease	Infusion, powder
Cough, fever, lean of blood, recovery of nervous, epilepsy, hysteria	Infusion
Cough, serious lung disease, skin disorders	Infusion, maceration in honey
Anemia, kidney and skin, cough, gall and liver disorders, better urinate, hepatitis, fever	Fresh juice, infusion, decoction
Hemorrhoids, lung tuberculosis and serious wounds, cough, nose irrigation	Decoction in milk Infusion, powder Hemorrhoids
Cough, skin disorder, liver disorders	Infusion, extract in olive oil
Cough at children, skin disease, gas in stomach elimination, scarlet fever	Infusion, tincture
Epilepsy, heart hardship, nematode elimination at children, bleeding in uterus and lung	Infusion, decoction, tincture
Menstrual disorders, against high potency, for nervous balance, hysteria.	Maceration, infusion, tincture
	Infusion, decoction, powder

these plants along with their main features is given in Table 1. All recorded species belong to vascular plants. The most frequently used plants in the whole region are: *Thymus serpyllum* (32 times), *Crataegus monogyna* (31 times), *Salvia officinalis* (31 times), *Tilia cordata* (31 times), *Urtica dioica* (31 times), *Taraxacum officinale* (31), *Hypericum perforatum* (27 times) *Achillea millefolium* (26 times), *Thymus pulegioides* (23 times), *Artemisia absinthium* (21 times), *Teucrium montanum* (21 times), *Nepeta catharia* (17 times), *Orchis morio* (17 times) and others (Table 1). For the first time in BiH, the following species are recorded as ones used by ordinary people as medicinal herbs: *Dictamnus albus*, *Origanum heracleoticum*, *Pinus heldreichii*, *Psoralea bituminosa*, *Rubia peregrina*, *Teucrium polium* and *Vitex agnus-castus*. For the first time use of *Sedum sexangulare*, *Lilium cattaniae* and *Micromeria thymifolia* in Mediterranean area is

recorded. Preparations made of detected medicinal plants are used for the treatment of wide spectrum of human diseases. The majority of species (63 of them) are used for the treatment of different respiratory disorders (cough, inflammation of throat, various infections, malicious kind of illnesses etc.). Significant share of plants (75 species) is used for the treatment of chronic illnesses of urinary and genital system, most commonly inflammation of urinary paths, inflammatory processes in the kidneys, formation of stone in the kidneys and the urinary bladder, then fungal or bacterial infections and as a preventive measure against the sterility, especially in females, as well as for treatment and rising of male potency. About 55 of analyzed species are primary used for treatment of gastrointestinal illnesses (ulcers of stomach and duodenum, digestion disorders, regulation of stomach acidity, bettering of appetite etc.). Significant number of

species is traditionally used for treatment of heart malfunctions, especially rhythm disorders, myocardial ischemia and circulation disorders, as well as for the treatment of skin conditions, wounds, diseases of nervous system (youth madness, depression, and fear), liver and gall diseases and metabolic disorders (Figure 2).

Shoot system (stem and leaves) is the most frequently used in medicinal purposes (Table 2), where usage of dried floral parts (34%) and dried leaves (around 21%) is predominant. Share of use of root system (rhizome, root, bulb and tuber) is approximately around 17% (Table 2).

The most frequent preparations for medicinal treatments from plants are usually very simple where infusion comprise around 44%, then diversity of decocts (around 19%), fresh juices (around 8%), and others (macerations, oils, tincture, etc.) (Table 3).

In the sense of taxonomy, plants that are being

used for therapeutic purpose in this region belong to divisions of Pteridophyta (one species from Equisetaceae, Aspleniaceae and Adiantaceae) and Spermatophyta with its two subdivisions: Coniferophytina (one species from *Cupressaceae* and *Pinaceae*) and Magnoliophytina (with two classes Magnoliopsida and Liliopsida, including 35 families all together). Domination of Magnoliopsida has been very noticeable with its 32 families comprised of 84 species that are used in therapeutic purposes. Labiatae family (Lamiaceae) represents the highest portion 20, followed by Compositae (Asteraceae) 11, Rosaceae 7, Leguminosae (Fabaceae) 4 and Umbelliferae (Apiaceae) 4. Class Liliopsida is represented by five families, 2 species are Liliaceae and one species each from Asparagaceae, Convallariaceae, Araceae and Orchidaceae. The rest of the families are represented by one or two species only (Table 1 and 4).

According to the EUNIS classification (EUNIS, 2006), 22 different habitats have been detected in the area of interest and they can be divided into several groups: woods and shrubs, meadows and rock debris communities, rock crevices and scree, swamps and abandoned places (Figure 3). The majority of medicinal plants (40%) are being found by people in open places, such as Mediterranean and sub-Mediterranean rock debris communities from the orders *Thero-Brachypodietalia* and *Scorzonero-Chrysopogonetalia*, termophilous meadows from order *Brometalia erecti* and wet meadows from the orders *Arrhenatheretalia* and *Trifolio-Hordeetalia secalini*. About 30% of species grows in the evergreen woods belonging to the order *Quercetalia ilicis*, termophilous broadleaved woods and shrubs from the orders *Quercetalia pubescentis* and *Ostryo-Carpinetalia orientalis* and black pine woods from the order *Pinetalia heldreichii-nigrae*. Significant number of species (about 20%) inhabits abandoned places (Figure 3).

DISCUSSION

Through the analysis of records of plant based medicinal treatments (Table 1), it becomes obvious that some of the plants are being used more frequently than the others (*Achillea millefolium*, *Hypericum perforatum*, *Salvia officinalis*, *Taraxacum officinale*, *Thymus serpyllum*, *Tilia cordata*, and *Urtica dioica*). These species were the most quoted by the informants in the entire area of investigation. Besides this, majority of these species play an important role in therapy of other regions, too (Antonone et al., 1988; Merzouki et al., 2000; Agelet and Valles's, 2001, 2003a, b; Bnouham et al., 2003; Loi et al., 2004; Gülcin et al., 2004; Novais et al., 2004; Guarrera et al., 2005a, b; Daher et al., 2006). Some of these species are used in a different way than in any other country. One such case is *Salvia officinalis* that in majority of Mediterranean countries is being used for throat rinsing,

after the tooth extraction and wound healing, while in the investigated area it is used to treat, first of all, heart diseases and then nervous system illnesses and kidney and liver problems (Lalićević and Djordjevic, 2004; Edi et al., 2005). This is also true for *Thymus serpyllum*. While the people of other areas use it as an excellent bronchial dilators (Rasoli and Mirmostafa, 2002), in this region it is used to treat heart diseases, hand tremor or to regulate the period. Species with wide distribution *Urtica dioica* is usually applied as a mean against anemia and cardiovascular diseases (Testai et al., 2002), but people in the area of interest also use it against a cough, fever, liver and gall diseases. In the entire world, *Taraxacum officinale* is known as a plant with the underlined healing character and here it is also used to treat anemia, hemorrhoids and jaundice. *Hypericum perforatum* is being used to treat different skin conditions in many countries, but in the area of investigation, its main purpose is to heal lung diseases and inflammation of kidneys. Next good example represent usage of *Tilia cordata* and *Tilia argentea* inflorescence that are used to treat illnesses of nervous system, such as epilepsy and hysteria, while in other parts of the world they are used as a medicine against a cough, mean to provoke sweating and in recent time for anti inflammatory effects (Anesini et al., 1999; Toker et al., 2004).

Comparative analysis of the ethno botanical practice between investigated area and other territories in the region (Capasso et al., 1982; De Feo et al., 1991; 1992; 1993; Malamas and Marselos, 1992; Bonet et al., 1999; Ballero et al., 2001; Comejo-Rodríguez et al., 2003; Guarrera, 2005; Ghorboni, 2005; Everest and Ozturh, 2005; Azaizeh et al., 2006; Avci et al., 2006) has shown that some identified plants possess many healing properties as they are being used to treat different diseases that are often diseases of completely different physiological systems. Thus, there is a group of plants that locals call "plants used for the treatment of any sickness" that is comprised of following species: *Achillea millefolium*, *Acorus calamus*, *Althaea officinalis*, *Arctium lappa*, *Artemisia absinthium*, *Centaurium maritimum*, *Chelidonium majus*, *Equisetum arvense*, *Hypericum perforatum*, *Juniperus oxycedrus*, *Juglans regia*, *Pimpinella saxifraga*, *Polygonum aviculare*, *Salvia officinalis*, *Sambucus ebulus*, *Thymus serpyllum* (including several species of the same genus) and *Teucrium montanum*. The good example is *Arctium lappa* that is used to treat skin conditions, to strengthen the hair root, against intestinal parasites, for dissolution of kidney stone, for improved urination, against diabetes and rabies, dog's bites, to release stomach gases, against sexually transmitted diseases, to treat facial nerve inflammation etc. Leaves of walnut *Juglans regia*, in addition to common healing the wounds are also used for the treatment of bacterial infections of female reproductive organs, liver diseases, various inflammations, strengthening the muscles and enlargement of the female




Figure 2. The frequency of wild medicinal plants used in human therapy. 1. Respiratory system, 2. Skin and hair, 3. Gastrointestinal system, 4. Genital system, 5. Liver, 6. Urinary system, 7. Miscellaneous, 8. Nervous system, 9. Gall disorders, 10. Blood system, 11. Heart disorders, 12. Bone and muscles, 13. Mouth, nose and throat, 14. Pancreas, 15. Spleen, 16. Bites of animals, 17. Eyes.

Table 2. Plant parts used in preparation.

Plant part	Number of plant part	Proportion (%)
Aerial part (all plant with flower)	42	34.15
Leaf	26	21.14
Fresh leaf	3	2.44
Young shoots	2	1.63
Flower	8	6.50
Fruit	9	7.32
Immature fruit	1	0.81
Seed	3	2.44
Rhizome	5	4.06
Root	14	11.38
Tuber and bulb	2	2.08
Bark	5	4.06
Resin	2	1.62
Fresh juice	1	0.81
Total:	123	100

breasts, raising a male potency, etc.

It has also been noticed that some medicinal plants, which are very popular in the Mediterranean area and wider, are rarely used in human therapy of population from the area of interest, such as: *Ruscus aculeatus*, *Punica granatum*, *Verbascum thapsus*, *Fraxinus ornus*, *Glycyrrhiza glabra*, *Herniaria hirsuta*, *Paliurus spina-*

christi, *Cnicus benedictus* and *Vitex agnus-castus*. Medicinal plants of the investigated area are being used to prevent and heal 140 different "diseases". In the course of this research, 527 ways of herbal usage in human phytotherapy have been identified (Table 1 and Figure 2). It is common that the same species is used for the treatment and prevention of several different

Table 3. Preparations.

Type of preparation	Number of species	Proportion (%)
Infusion	78	41.71
Light infusion	1	0.53
Infusion in vinegar	1	0.53
Infusion with honey	1	0.53
Infusion in cooked wine	1	0.53
Extract in Olive oil	3	1.60
Extract in wine	2	1.07
Decoction	23	16.67
Decoction in milk	3	1.60
Decoction in wine	1	0.53
Beverage	1	0.53
Fresh fruit	2	1.07
Fresh juice	14	8.33
Maceration	9	4.17
Maceration in honey	1	0.53
Powder	13	2.08
Resin	2	2.08
Tincture	25	1.04
Oil	2	1.04
Traditional balm	2	1.07
Fresh crushed leaves	1	0.53
Crushed leaves with honey	1	0.53
Total:	187	100

Table 4. Taxonomy belonging of medicinal plants to botanical family.

No	Plant family	Number of species	Proportion (%)
1	<i>Lamiaceae</i>	20	20.83
2	<i>Asteraceae</i>	11	11.46
3	<i>Rosaceae</i>	7	7.29
4	<i>Fabaceae</i>	4	4.17
5	<i>Apiaceae</i>	3	3.12
6	<i>Crassulaceae</i>	3	3.12
7	<i>Liliaceae</i>	2	2.08
8	<i>Malvaceae</i>	2	2.08
9	<i>Papaveraceae</i>	2	2.08
10	<i>Anacardiaceae</i>	2	2.08
11	<i>Boraginaceae</i>	2	2.08
12	<i>Rutaceae</i>	2	2.08
13	<i>Rhamnaceae</i>	2	2.08
14	<i>Pinaceae</i>	2	2.08
15	<i>Violaceae</i>	2	2.08
16	The other 29 families have been presented by one plant species	31	32.29
	Total:	96	100

diseases as its healing properties influences different functional systems. Thus the study has shown that the average number of health disorders being treated by the

same plant is approximately 4 - 5. For example *Arctium lappa* is applied in 12 cases, *Achillea millefolium* in 11, *Salvia officinalis* in 9, *Polygonum aviculare* and *Juglans*




Figure 3. Wild medicinal plants in relate to their habitat and community. 1. Forest and scrubs; 2. Grassland and rocky grasslands; 3. Wetlands; 4. Cliffs and scree; 5. Abandoned habitats.

regia in 8, *Sambucus ebulus*, *Pimpinella saxifraga* and *Centaureum maritimum* in 7 etc. Most of the plants are being used for medicinal treatment of five different illnesses. There are only few species which usage is limited to only one, two or three diseases (*Adiantum capillus-veneris*, *Eryngium campestre*, *Geranium macrorrhizum*, *Lilium cattaniae*, *Pinus nigra*, *Pistacia lentiscus*, *Polygonum odoratum*, *Polypodium vulgare*, *Punica granatum*, *Sedum maximum*, *Sedum sexangulare*). Some plants treat disorders of a single functional system, such as entire genus *Teucrium* (*T. montanum*, *T. chamaedrys*, *T. arduini*, and *T. polium*) that is only used for the treatment of digestive system.

Through the comparative record's analysis (Table 1 and Figure 2), one comes to the conclusion that despite being uneducated or lacking basic medicinal knowledge, common people make a positive difference in the treatments of numerous unfavorable health conditions (over 140). Majority of species are being used to fight respiratory system difficulties (63), especially against cough (even 32 species) and asthma and lung diseases (16). There are few species, such as *Chelidonium majus*, *Sedum acre* and *Satureja subspicata* that are commonly used to treat tuberculosis or lung cancer. Digestive system is often an object of herbal medicinal treatment too and over 30 different diseases or disorders are prevented and cured by herbal preparations. 55 species are used for stomach and intestinal therapy of which major proportion is for the stomach difficulties

(19 species) and digestion disorders on such as constipation and diarrhea (11 species). 41 species are being used in the treatment of liver sicknesses, including jaundice and the gall. Treatment comprised of five plants has achieved a great success in the maintenance of stable sugar level in the bloodstream.

Traditional phytotherapy provides excellent results in the treatment of diverse cardiovascular disorders, especially arrhythmia, heart fibrillation and the diseases of blood vessels. In the investigated area, unlike other parts of BiH (Redzic, 2006b) or wider (Ivancheva and Stantcheva, 2000; Pieroni et al., 2003; Jaric et al., 2007), different heart disorders have been cured with great success. For this, preparations of *Satureja montana* and *S. subspicata* have been used, while in other regions *Artemisia absinthium* is used for the same purpose, as well as for treatment of arrhythmia. *Acorus calamus* is used for the treatment of heart fibrillation while *Salvia officinalis* is well known in the investigated area as a „cardiac treatment plant“ used to strengthen heart muscle and to treat arrhythmia. *Thymus serpyllum* strengthens heart muscle, etc. There are many species that regulate nervous system and by that positively regulate cardiovascular system too.

There are several plants used for the treatment of diseases that affect blood, mainly in terms of «blood cleansing» and against anemia, while some of them, such as *Satureja montana* and local endemic species *Satureja subspicata*, are used to treat leukemia and

syndromes of lymphatic nodes since long ago. There is an interesting story of a 70 years old man from nearby Stolac, who was as a 23 old boy dismissed from the military academy in Belgrade for his bad health, with a life expectancy of only 2 - 3 months. Afterwards, he was subjected to the intense therapy of powder and infusion prepared with *Satureja montana* and *S. subspicata* (combined with goat's milk). The final result had astonished doctors in the region and he started his new life as a healthy man where medicinal plants became an obligatory part of it.

Thirty-three plants are applied in the treatment of kidney and urinary system diseases, mainly as a mean to dissolve a stone in kidneys, for the treatment of urinary infections and inflammation processes, or to improve urination. The best effects have been reached by following species: *Arbutus unedo*, *Adiantum capillus-veneris*, *Arctium lappa*, *Asparagus officinalis*, *Carlina acaulis*, *Hieracium pilosella*, *Hypericum perforatum*, *Polygonum aviculare*, *Sambucus ebulus* and others.

At present, plants are being used for the treatment of kidney and urinary system diseases in other regions too (Abu-Rabia, 2005; Lans, 2006). 42 plant species have been used in the treatments of conditions affecting genital system. Informants have suggested 21 different modes of usage of these plants, covering all possible unfavorable conditions of genital system. Majority of plants (13 species) are used for the healing of female genital infections (*Achillea millefolium*, *Artemisia vulgaris*, *Capsella bursa-pastoris*, *Juglans regia*, *Pimpinella saxifraga*, *Plantago meda*, *Quercus pubescens*, *Potentilla anserina*, *Mentha pulegium*, and especially *Sedum sexangulare*). 5 species have been used for the treatment of prostate inflammatory processes (*Asparagus officinalis*, *Hieracium pilosella* and especially *Nigella damascena*). The underlined use of plants as an aphrodisiac aimed to raise male potency (*Juglans regia*, *Satureja montana*, *Hyssopus officinalis*, especially *Orchis morio* and *Origanum heracleoticum*) reflects southern temperament of the people in the investigated area. Significant experiences in the usage of traditional herbal remedies for female gynecological disorders have also been recorded in the other parts of the world (Arnel-Schnebelen et al., 2004; Vidyasagar and Prashantkumar, 2007).

Fifty-six species have been used for the treatment of different skin conditions of which the most significant are: *Althaea officinalis*, *Arctium lappa*, *Carlina acaulis*, *Cotinus coggygria*, *Pinus nigra*, *Pistacia lentiscus*, *Polypodium vulgare*, *Sedum maximum* and *Viola alba*. Emphasized use of medicinal plants for the treatment of skin conditions has been recognized in some other regions as well (Burnstock et al., 1999; Onanga et al., 1999; Saikia et al., 2006).

Significant results are also being reached in terms of treatments of nervous system disorders such as hysteria, epilepsy and psychotic state of mind for which 29 plant

species have been used. The most efficient are: *Hypericum perforatum*, *Acorus calamus*, *Vitex agnus-castus*, *Dictamnus albus*, *Glechoma hederacea*, *Micromeria thymifolia*, *Nepeta cataria*, *Mentha pulegium*, *Thymus pulegioides*, and *Viscum album*.

About 20 plants are used to treat the diseases of muscles and skeletal system, mouth, nose and throat (Table 1). It should be emphasized that *Sanguisorba minor* is used to treat bites of the most venomous snakes in the area, such as *Vipera berus* and *Vipera ammodytes*.

This characteristic of *Sanguisorba minor* arises from its content of unique phenol carboxylic acids (Nahla, 2003). In the region of high biological diversity, where poisonous animal bites occur, seeking for a plant that can treat these states is a specially protruding (Otero et al., 2000).

Similar to other countries (Bandeira et al., 2001; Palmese et al., 2001; Passalacqua et al., 2007) in terms of plant's part being used, the most prominent practice is use of aerial parts where usage of flowering plants (about 34%) and leaves (about 22%) dominate. This proportion correlates with the general domination of hemi cryptophytes in the area (meaning perennial plants), as well as the simple way of collecting them. This isn't the case with trees or plants developing massive root system organs, making a collection more complicated, in a karsts dominated territory (substratum is very hard to penetrate). Even though the proportion of geophytes in the investigated area is much higher, the usage of the root system (root, bulb and rhizome) for the medicinal treatments comes from only 19 plants. Similar to the other territories in BiH (Redzic et al., 1989; 1991; 2006b), usage of coniferous tree's resin as an important ingredient of some traditional pharmaceuticals is very popular here. Pharmaceutical preparations are often prepared from several species of plants, (usually mixing three, five or up to seven). It is believed that the effect of cure is better if more plants are used in its preparation.

Majority of plant parts, which are usually aerial parts, are being used to prepare diverse infusion. These preparations are made with simple boiling of plant leaves or flowers in the water, removing it from a heat and leaving it covered until cooled. Afterwards, the infusion is filtrated and served in a traditional manner with the sage honey. A cup of this should be taken several times a day. It can also be consumed instead of water or any other liquid, such as infusion of *Teucrium montanum* (usually 2 - 3 L of tea are made in one go). Jaundice is very successfully treated in this way, as well as liver and stomach diseases, depression etc. Maceration is prepared in way where a crushed plant has been sunk in the warm water over the night.

Homemade tinctures, also frequently used for treatments, are prepared in the way that plant parts are sunk into brandy, kept on the sunlight for 7 - 10 days and applied in a form of drink or cataplasm to cure stomach aches, respiratory system diseases and skin conditions.

Unique preparations are creams known as "melehem"

whose traditional use in this area is very long. They are prepared using fresh plant parts or pine's resin and mixed with cow's butter or pig's fat. *Mehlems* are common cure for hard healing wounds.

Apart from home made medicinal preparations, medicinal plants are commonly used for a traditional nutrition. The dishes are mainly made with olive oil (Redzic, 2006a).

Compared to the international databases on medicinal and edible plants (PFAF, 2006), and to the other countries in Mediterranean region (Ivancheva and Stantcheva, 2000; Pieroni et al., 2002b; 2003; El-Hilaly et al., 2003; Pieroni and Quave, 2005; Scherrer et al., 2005) to the area of interest, significant difference regarding plant diversity has been detected, including its taxonomical position, ecological character, way of preparing medicinal substances and forms of its application in traditional human therapy.

Comparison of obtained results to the most encompassing database plants for the future (PFAF, 2006), has lead to discovery of 17 species whose therapeutic usage was mentioned for the first time. This represents an important contribution to the advancement of ethno botany and pharmaceuticals on global scale (Table 1). Special attention, in terms of being valuable therapeutic resource, should be paid to: *Teucrium montanum* (that according to local people beliefs «heal every disease»), *Teucrium arduini* (endemic species), *Centaureum maritimum* (very popular plant when it comes to stomach and «female» sickness) and *Eryngium amethystinum* (used to treat «water sickness» and hepatitis). Despite being SE European endemic species, *Eryngium amethystinum* posses wide distribution in the investigated area, representing therefore secure resource for further research and modern therapeutic practice.

Using the above mentioned database, it appears that *Helichrysum italicum* hasn't been known for its medicinal characteristics so far and it was only recorded as a spice. Nevertheless, in the area of interest, this is the mostly used medicinal plant used in traditional therapy for liver, gall and stomach sickness, plus it works as an excellent insecticide. The list of species with so far less known or even completely unknown therapeutic use includes the following: *Lilium cattaniae* (local endemic plants), *Marrubium incanum* (excellent anti-arrhythmic), *Micromeria thymifolia* (has an odor similar to that of genus *Mentha* and *Thymus*), *Origanum heracleoticum* (has a luscious smell), resin of the endemic kind of pine *Pinus heldreichii*, then *Prunella laciniata*, *Psoralea bituminosa* (used in the treatment of skin conditions), *Rubus heteromorphus* (used in the treatment of diarrhea and cough) and *Salvia bertolonii* (used in the treatment of skin conditions and cough). Comparing to the same database, *Sanguisorba minor* has been used rarely in the therapy and yet, in the area of interest, this plant is used to treat bites of the most venomous snakes that frequently occur in this area, such as *Vipera ammodytes*

and *Vipera berus* subsp. *bosniaca*. It is also used for the treatment of diabetes.

Species such as *Satureja subspicata* (used to treat the most severe blood diseases, such as leukemia), *Sedum maximum* and *Sedum sexangulare* (used to treat skin conditions and complicated vaginal infections) have also been unknown to the world's ethno botanical databases: Further ethno pharmacological research is needed to confirm true potential of these plants in medicinal treatment in the above named or any other pathological condition in human health.

The comparison of the generated data with the international databases in the pertaining field (PFAF, 2006) and mentioned references revealed that the species of *Centaureum maritimum* (L.) Fritsch (Syn: *Centaureum discolor* (Gand.) Ronniger) have not been registered previously to have any use in the traditional therapy. Comparative studies of this plant to the one preformed on very popular species in ethno botany *Centaureum umbellatum* (Grujic-Vasic et al., 2005) showed that the species *C. maritimum* have a significantly higher degree of bitterness. This may indicate its potentially higher phytochemical capacities, and should be confirmed in a concrete pharmacological research. The traditional tincture of „grape brandy“ with an aerial part of the species *C. maritimum* has been proved to have an excellent influence on heart strengthening as well as on elimination of stomach discomfort. It is also very popular as an everyday aperitif.

There is no previous record of species *Eryngium amethystinum* L. (Syn.: *Eryngium multifidum* Sibth. and Sm.; *Eryngium glomeratum* Lam.) in the human therapy. This generates the need for more detailed both phytochemical and bio-assay research efforts aimed at this productive species, because complex researches of similar species of the genus *Eryngium* (in particular *Eryngium campestre*, *Eryngium maritimum*) indicate obvious anti-inflammatory activity (Kupeli et al., 2006). As a matter of fact, some species of this genus such as the more widely distributed *Eryngium campestre* contains cyclohexenone and cyclohexadienone glycoside (Clemens et al., 1986).

So far, species *Marrubium incanum* Desr. (Syn.: *Marrubium candidissimum* auct., non L.) was not registered in human therapy even though, according to this research, has shown to be very effective in the treatment of not only stomach diseases but also heart conditions such as arrhythmia. It is extremely important to identify pharmacological effects of this species, the more so because a seed of *Marrubium candidissimum* contains anti Tn (Bird and Wingham, 1981).

Lilium cattaniae (Vis.) Vis. (Syn.: *Lilium martagon* L. var. *cattaniae* Vis.; *Lilium. martagon* L. var. *atropurpureum* Neilr.; *Lilium dalmaticum* Vis. in Sched.) is an endemic, very decorative species. It grows in thermophilous forests and shrubs of Herzegovina. So far, it was neither recorded in traditional therapy, nor it was

researched from the phytochemical aspect. It represents an important resource both for phytochemical and pharmacological research, because its closest species *Lilium martagon* in the neighboring Albania is being used in therapy „to eliminate lipids from the blood“, to treat every liver disease (Pieroni et al., 2005b), and it also contains pyrrolidine glycoside ester and steroidal saponins (Satou et al., 1996).

Origanum heracleoticum L. (Syn.: *Origanum hirtum* Link) is an endemic species of strong odor. So far, it was not recorded in the human therapy in the Mediterranean region. The research that was conducted on this species showed that its essential oil has a significant antimicrobial activity on *Pseudomonas putida* (Oussalah et al., 2006), *Escherichia coli* O157:H7, *Salmonella Typhimurium*, *Staphylococcus aureus* and *Listeria monocytogenes* (Oussalah et al., 2007). It also contains very unique chemical compositions such as 4, 5-epoxy-p-menth-1-ene (Lawrence et al., 1974), and a series of other organic compositions (Nhu-Trang et al., 2006). This indicates a need for thorough bio-assay research of these very popular species in BiH coastline.

Pinus heldreichii H. Christ (Syn.: *Pinus leucodermis* Antoine p.p.), endemic Balkan-Apennine pine with its distribution in the Central and Western Balkans, is used in the human therapy in the area of interest. Its usage in the other areas has not been recorded. Tree sap is mainly used from this species. This may indicate a medicinal value of both twigs and bark because it contains cembratrienols and other organic compounds at white bark of this species (Werner et al., 1994).

Species *Prunella laciniata* (L.) L. (Syn.: *Prunella alba* Pall. ex M. Bieb.), which mainly inhabits warmer habitats of open sub-Mediterranean meadows, so far hasn't been recorded in human phytotherapy either. In the surveyed area, this species is used for making infusions for the cough treatment, gall and skin diseases. Its closest relative species, the more widely distributed *Prunella laciniata*, is used externally „for cleaning open sores and wounds“ (Rivera and Obon, 1995). Given that *Prunella laciniata* inhabits ecologically clean habitats and it is numerous, it could be considered as a serious resource in modern phytotherapy.

Bitumen trefoil *Psoralea bituminosa* L. is Mediterranean species which so far has not been used by humans in BiH. However, it was recorded in therapy of Madeira and Porto Santo, where they use its leaves for making „decoction with alcohol and iodine that is applied externally as a hair restorer“ (Rivera and Obon, 1995).

For the first time, *Salvia bertoloni* Vis. (Syn.: *Salvia pratensis* L. subsp. *bertoloni* (L.) Visiani) has been mentioned in human therapy of Mediterranean countries where this species is located. There are several indicators that this is truly a medicinal plant because its closest relative, a typical meadow sage *Salvia pratensis* L. in addition to a series of compositions which have an antioxidant effect (Miliauskas et al., 2004; Lima et al.,

2005; Eidi et al., 2005), contains lupeol derivatives (Anaya et al., 1989). It is also necessary to make a thorough investigation of pharmacological effects of the species *Salvia bertoloni*, which may have been unjustifiably left out in favor of the more popular medicinal plant *Salvia officinalis*, since the two frequently share same habitats.

Red heather *Satureja subspicata* (Syn.: *Satureja illyrica* Host; *Satureja montana* L. subsp. *illyrica* Nyman) is a Balkan endemic species with its center of distribution in the investigated area (Silic, 1984). So far, its official usage in a therapy has not been recorded. It was probably pushed out from the use by the species *Satureja montana*, which is more widely distributed and more popular among people. Having in mind an extremely prominent chemical diversity and antimicrobial activity of the species *Satureja subspicata* from the habitat of the neighboring Croatia (Skočibušić et al., 2006), it is necessary to do more detailed pharmacological researches, because of a belief that this species successfully treats and „protects“ from very serious diseases.

Species of the genus *Sedum* such as *Sedum sexangulare* (Syn.: *Sedum mite* auct.; *Sedum boloniense* Loisel.), as well as *Sedum maximum* (L.) Suter (Syn.: *Sedum telephium* L. subsp. *maximum* (L.) Krock) have not been registered in the therapy of either BiH region or in the wider Mediterranean region. Having in mind their role in treating skin diseases, infections, vaginal discharge and liver, it would be necessary to complete pharmacological researches on these species. The reason for this is mostly because of flavonol glycosides (Mulinacci et al., 1995) that were isolated in the leaves of *Sedum maximum*, which have been used to treat wounds during the last war in BiH.

Although the popular *Teucrium montanum* L. (Syn.: *Teucrium pannonicum* A. Kern.; *Teucrium praemontanum* Klokov; *Teucrium helianthemoides* Adamovic; *Teucrium jailae* Juz.) has been in use for a long time, its effects are yet to be sufficiently pharmacologically researched and its medicinal effects confirmed. Available phytochemical researches pointed at the presence of neo-clerodane diterpenoids (Malakov et al., 1992), and the populations of this species from the region of the neighboring Serbia, has indicated the antimicrobial and free radical scavenging activities (Djilas et al., 2006). In particular, it would be necessary to examine its pharmacological effects on the human liver.

Dinaric endemic species *Teucrium arduinii* L. (Syn.: *Teucrium hircanicum* auct. eur., non L.) is a sort of novelty in the therapy but it deserves a full attention in the sense of both more detailed phytochemical and pharmacological researches, because of neo-cleodane diterpenoids indication of an anti-feedant activity (Bruno et al., 2002).

True significance of possible usage of the abovementioned wild plants can be revealed through modern tendencies in the ethno pharmacology (Gilani

and Rahman, 2005) and through the stronger connection between the ethno botany and ethno pharmacology (Heinrich et al., 2006).

ACKNOWLEDGEMENTS

This paper was co-financed by means from the scientific-investigation project: “*Procjena ugroženosti i obrasci održive zaštite genofonda endema u razvojnom centru Prenj-Čvrstica-Čabulja*” (“Assessment of threats and patterns for the sustainable protection of endemic gene pool in the development centre Prenj-Čvrstica-Čabulja”), Project Number: 03-39-5980-67-2/08, co-financed by the Federal Ministry of Education and Science of Federation of Bosnia and Herzegovina.

The author owes his gratitude to the people who have supported me during the challenging field investigation, especially to Mr. Sedik Velic, technician of the Dep. of Botany of the Faculty of Science University of Sarajevo. I am also very thankful to the English language experts Ms. Sabina Trakic, Mr. Mirnes Zukanovic and Miss. Fadila Muftic, for their efforts made during the translation of this paper to English, and the entire revision of original text written in Bosnian language.

REFERENCES

- Abella L, Cortella AR, Velasco-Negueruela A, Pérez-Alonso MJ (2000). Ethnobotany, volatile oils and secretion tissues of *Werneria poposa* from Argentina. *Pharm. Biol.* 38: 197-203.
- Abu-Rabia B (2005). Urinary diseases and ethnobotany among pastoral nomads in the Middle East. *J. Ethnobiol. Ethnomed.* 1: 4.
- Agelet A, Valle's J (2001). Studies on pharmaceutical ethnobotany in the region of Pallars (Pyrenees, Catalonia, Iberian Peninsula). Part I. General results and new or very rare medicinal plants. *J. Ethnopharmacol.* 77: 57-70.
- Agelet A, Valles J (2003a). Studies on pharmaceutical ethnobotany in the region of Pallars (Pyrenees, Catalonia, Iberian Peninsula). Part II. New or very rare uses of previously known medicinal plants. *J. Ethnopharmacol.* 84: 211-227.
- Agelet A, Valles J (2003b). Studies on pharmaceutical ethnobotany in the region of Pallars (Pyrenees, Catalonia, Iberian Peninsula). Part III. Medicinal uses of non-vascular plants. *J. Ethnopharmacol.* 84: 229-234.
- Anaya J, Caballero MC, Grande M, Navarro JJ, Tapia I, Almeida JF (1989). A lupeol derivative from *Salvia pratensis*. *Phytochem.* 28: 2206-2208.
- Anesini C, Werner S, Borda E (1999). Effect of *Tilia cordata* flower on lymphocyte proliferation: participation of peripheral type benzodiazepine binding sites. *Fitoterapia* 70: 361-367.
- Antonone R, De Simone F, Morrica P, Ramundo E (1988). Traditional phytotherapy in the Roccamonfina volcanic group, Campania, Southern Italy. *J. Ethnopharmacol.* 22: 295-306.
- Applebeck WL (2004). Medicinal plants in folk tradition: An ethnobotany of Britain and Ireland by David E. Allen and Gabrielle Hatfield. *Syst. Bot.* 29: 1021-1021.
- Arnel-Schnebelen B, Hadji-Minaglou F, Peroteau JF, Ribeyre F, de Billerbeck VG (2004). Essential oils in infectious gynaecological disease: a statistical study of 658 cases. *Int. J. Aromather.* 14: 192-197.
- Avci G, Kupeli E, Eryavuz A, Yesilada E, Kucukkurt I (2006). Antihypercholesterolaemic and antioxidant activity assessment of some plants used as remedy in Turkish folk medicine. *J. Ethnopharmacol.* 107: 418-423.
- Azaizeh H, Saad B, Khalil K, Said O (2006). The state of the art of traditional Arab herbal medicine in the Eastern Region of the Mediterranean: A Review. *Evid Based Complement Alternat. Med.* 3: 229-235.
- Bailly F, Queffelec C, Mbemba G, Mouscadet JF, Cotellet P (2005). Synthesis and HIV-1 integrase inhibitory activities of caffeic acid dimers derived from *Salvia officinalis*. *Bioorg. Med. Chem. Lett.* 15: 5053-5056.
- Ballero M, Poli F, Sacchetti G, Loi MC (2001). Ethnobotanical research in the territory of Fluminimaggiore (South-western Sardinia). *Fitoterapia* 72: 788-800.
- Bandeira SO, Gaspar F, Pagula FP (2001). African ethnobotany and healthcare: emphasis on Mozambique. *Pharm. Biol.* 39(1): 70-73.
- Bedoya LM, Sanchez-Palomino S, Abad MJ, Bermejo P, Alcami J (2001). Anti-HIV activity of medicinal plant extracts. *J. Ethnopharmacol.* 77: 113-116.
- Begovic B (1960). Strani kapital u sumskoj privredi Bosne i Hercegovine za vrijeme otomanske vladavine./ Foreign capital in the forest economy of BiH during the Ottoman empery. *Radovi Sumarskog fakulteta i Instituta zaumarstvo i drvnu industriju u Sarajevu/Papers of Faculty of Forestry and Institute for forestry and wood industry.* (In Serbo-Croatian). Sarajevo 5: 3-272.
- Bird GWG, Wingham J (1981). Anti-Tn from *Marrubium candidissimum*. *Rev. Fr. Transfus. Immunohematol.* 24: 347-348.
- Bnouham M, Merhfour FZ, Abderrahim Z, Mekhfi H, Aziz M, Legssyer A (2003). Antihyperglycemic activity of the aqueous extract of *Urtica dioica*. *Fitoterapia* 74: 677-681.
- Bodeker C, Ong CK, Grundy C, Burford G, Shein K (2005). WHO Global atlas of traditional, complementary and alternative medicine. World Health Organization, Center for Health Development, Kobe, Japan.
- Bonet MA, Parada M, Selga A, Valle's J (1999). Studies on pharmaceutical ethnobotany in the regions of L'Alt Emporda' and Les Guilleries (Catalonia, Iberian Peninsula). *J. Ethnopharmacol.* 68: 145-168.
- Bonet MÀ, Vallès J (2003). Pharmaceutical ethnobotany in the Montseny biosphere reserve (Catalonia, Iberian Peninsula). General results and new or rarely reported medicinal plants. *J. Pharm. Pharmacol.* 55: 259-270.
- Bruno M, Piozzi F, Maggio AM, Rosselli S, Simmonds MSJ, Servetaz O (2002). Antifeedant activity of neo-clerodane diterpenoids from *Teucrium arduini*. *Biochem. Sys. Ecol.* 30: 595-599.
- Burnstock G, Onanga M, Ekouya A, Ouabonzi A, Itoua GB (1999). Ethnobotanical, pharmacological and chemical studies of plants used in the treatment of 'Mwandza' dermatitis. *Fitoterapia* 70: 579-585.
- Capasso F, De Simone F, Senatore F (1982). Traditional phytotherapy in the Agri Valley, Lucania, Southern Italy. *J. Ethnopharmacol.* 6: 243-250.
- Carai MAM, Agabio R, Bombardelli E, Bourov J, Luigi Gessa G, Lobina C, Morazzoni P, Pani M, Reali R, Vacca G, Colombo G (2000). Potential use of medicinal plants in the treatment of alcoholism. *Fitoterapia* 71: S38-S42.
- Clemens A, Erdelmeier J, Sticher O (1986). A cyclohexenone and a cyclohexadienone glycoside from *Eryngium campestre*. *Phytochem.* 25: 741-743.
- Comejo-Rodríguez J, Ascensao L, Angels Bonet M, Valles J (2003). An ethnobotanical study of medicinal and aromatic plants in the Natural Park of Serra de Sao Mamede" (Portugal). *J. Ethnopharmacol.* 89: 199-209.
- Daher CF, Baroody KG, Baroody GM (2006). Effect of *Urtica dioica* extract intake upon blood lipid profile in the rats. *Fitoterapia* 77: 183-188.
- De Feo V, Ambrosio C, Senatore F (1991). Traditional phytotherapy in Caserta province, Campania, Southern Italy. *J. Ethnopharmacol.* 63: 337-349.
- De Feo V, Aquino R, Menghini A, Ramundo E, Senatore F (1992). Traditional phytotherapy in the Peninsula Sorrentina, Campania, Southern Italy. *J. Ethnopharmacol.* 36: 113-125.
- De Feo V, Senatore F (1993). Medicinal plants and phytotherapy in the Amalfitan Coast, Campania, Southern Italy. *J. Ethnopharmacol.* 39: 39-51.

- Djilas SM, Markov SL, Cvetkovic DD, Canadanovic-Brunet JM, Cetkovic GS, Tumbas VT (2006). Antimicrobial and free radical scavenging activities of *Teucrium montanum*. *Fitoterapia* 77: 401-403.
- Donaldson J, Rex C (2004): Screening for Anticancer Agents from Sonoran Desert Plants: A Chemical Ecology Approach. *Pharm. Biol.* 42: 478-487.
- Eidi M, Eidi A, Zamanizadeh H (2005). Effect of *Salvia officinalis* L. leaves on serum glucose and insulin in healthy and streptozotocin-induced diabetic rats. *J. Ethnopharmacol.* 100: 310-313.
- EI-Hilaly J, Hmammouchi M, Lyoussi B (2003). Ethnobotanical studies and economic evaluation of medicinal plants in Taounate province (Northern Morocco). *J. Ethnopharmacol.* 86: 149-158.
- EUNIS (2006). EUNIS Habitat Classification. (Available on line: <http://eunis.eea.eu.int/habitats.jsp>; Access 20. December 2006). European Environment Agency, Kongens Nytorv 6, DK-1050 Copenhagen K, Denmark.
- EURO-MED Plant Base (2006). The information resource for Euro-Mediterranean plant diversity. Page editor: G. Domina. This page last updated June 14, 2006. Dipartimento di Scienze botaniche ed Orto botanico, Università degli Studi di Palermo. (Available on line <http://www.emplantbase.org/home.html>, Accessed 20. December 2006).
- Everest A, Ozturk E (2005). Focusing on the ethnobotanical uses of plants in Mersin and Adana provinces (Turkey). *J. Ethnobiol. Ethnomed.* 1: 6.
- Fernandez EC, Sandi YE, Kokoska L (2003). Ethnobotanical inventory of medicinal plants used in the Bustillo Province of the Potosi Department, Bolivia. *Fitoterapia* 74: 407-416.
- Fukarek P (1954). Istrazivanja flore i vegetacije Bosne i Hercegovine./Research of flora and vegetation of Bosnia and Herzegovina. (In Croatian). *God Biol inst Uni Sarajevo/ J. Biol. Inst. Univ. Sarajevo* 7: 111-168.
- Ghorboni A (2005). Studies on pharmaceutical ethnobotany in the region of Turkmen Sahara, north of Iran: (Part 1): General results. *J. Ethnopharmacol.* 102: 8-68.
- Giday M, Asfaw Z, Elmquist T, Woldu Z (2003). An ethnobotanical study of medicinal plants used by the Zay people in Ethiopia. *J. Ethnopharmacol.* 85: 43-52.
- Gilani AH, Rahman A (2005). Trends in ethnopharmacology. *J Ethnopharmacol* 100: 43-49.
- Girach RD, Ahmad A, Ahmad M (1998): Medicinal ethnobotany of Sundargarh, Orissa, India. *Pharm. Biol.* 36: 20-24.
- Gluck L (1892). Narodni lijekovi iz bilinstva u Bosni. *Glasnik Zem muz, Sarajevo* 4: 134-167.
- Grujic-Vasic J, Pilipovic S, Redzic S, Ibrulj A, Mijanovic M (2005). Phenolic acids and antiinflammatory activity of *Centaureum umbellatum* Gillib. (*Centaureum erythraea* Rafn. (Gentianaceae). Works of Academy of Science and Arts of BiH, Dep. of Med. Sci. 94(34): 3-11.
- Guarrera PM (2005). Traditional phytotherapy in Central Italy (Marche, Abruzzo, and Latium). *Fitoterapia* 76: 1-25.
- Guarrera PM, Salerno G, Caneva G (2005). Folk phytotherapeutical plants from Maratea area (Basilicata, Italy). *J. Ethnopharmacol.* 99: 367-378.
- Guarrera PM, Forti G, Marignoli S (2005b). Ethnobotanical and ethnomedicinal uses of plants in the district of Acquapendente (Latium, Central Italy). *J. Ethnopharmacol.* 96: 429-444.
- Gülcin I, İrfan Küfrevioğlu O, Oktay M, Büyükkuroğlu ME (2004). Antioxidant, antimicrobial, antiulcer and analgesic activities of nettle (*Urtica dioica* L.). *J. Ethnopharmacol.* 90: 205-215.
- Heinrich M (2003). Ethnobotany and natural products: the search for new molecules, new treatments of old diseases or a better understanding of indigenous cultures? *Curr. Top. Med. Chem.* 3: 141-154.
- Heinrich M (2005). Ethnobotany and natural products: the search for new molecules, new treatments of old diseases or a better understanding of indigenous cultures?. *Front Med. Chem. Online* 2: 431-450.
- Heinrich M, Bremner P (2006). Ethnobotany and Ethnopharmacy – Their Role for Anti-Cancer Drug Development. *Curr. Drug Targets* 7: 239-245.
- Heinrich M, Kufer J, Leonti M, Pardo-de-Santayana M (2006). Ethnobotany and ethnopharmacology—Interdisciplinary links with the historical sciences. *J. Ethnopharmacol.* 107:157-160.
- IUCN (1996). *Endangered Species Resources*. (Available on line (<http://www.usd320.k12.ks.us/whs/lmc/endangeredspecies.html>; Accessed 28. December, 2006).
- Ivancheva S, Stantcheva B (2000). Ethnobotanical inventory of medicinal plants in Bulgaria. *J. Ethnopharmacol.* 69: 195-172.
- Jaric S, Popović Z, Mačukanović-Jocić M, Djurdjević L, Mijatović M, Karadžić B, Mitrović M, Pavlović P (2007). An ethnobotanical study on the usage of wild medicinal herbs from Kopaonik Mountain (Central Serbia). *J. Ethnopharmacol.* 111: 160-175.
- Kenny MG (2002). A darker shade of green: medical botany, homeopathy, and cultural politics in interwar Germany. *Soc. Hist. Med.* 15: 481-504.
- Kupeli E, Kartal M, Aslan S, Yesilada E (2006). Comparative evaluation of the anti-inflammatory and antinociceptive activity of Turkish *Eryngium* species. *J. Ethnopharmacol.* 107: 32-37.
- Lalićević S, Djurdjević I (2004). Comparison of benzydamine hydrochloride and *Salvia officinalis* as an adjuvant local treatment to systemic nonsteroidal anti-inflammatory drug in controlling pain after tonsillectomy, adenoidectomy, or both: an open-label, single-blind, randomized clinical trial. *Curr. Ther. Res.* 65: 360-372.
- Lans CA (2006). Ethnomedicines used in Trinidad and Tobago for urinary problems and diabetes mellitus. *J. Ethnobiol. Ethnomed.* 2: 45.
- Lawrence BM, Terhune SJ, Hogg JW (1974). 4,5-Epoxy-*p*-menth-1-ene: A new constituent of *Origanum heracleoticum*. *Phytochem.* 13: 1012-1013.
- Leduc C, Coonishish J, Haddad P, Cuerrier A (2006). Plants used by the Cree Nation of Eeyou Istchee (Quebec, Canada) for the treatment of diabetes: A novel approach in quantitative ethnobotany. *J. Ethnopharmacol.* 105: 55-63.
- Leporatti ML, Corradi L (2001). Ethnopharmacobotanical remarks on the Province of Chieti town (Abruzzo, Central Italy). *J. Ethnopharmacol.* 74: 17-40.
- Leporatti ML, Ivancheva S (2003). Preliminary comparative analysis of medicinal plants used in the traditional medicine of Bulgaria and Italy. *J. Ethnopharmacol.* 87: 123-142.
- Lev E, Amar Z (2002). Ethnopharmacological survey of traditional drugs sold in the Kingdom of Jordan. *J. Ethnopharmacol.* 82: 131-145.
- Lima CF, Andrade PB, Seabra RM, Fernandes-Ferreira M, Pereira-Wilson C (2005). The drinking of a *Salvia officinalis* infusion improves liver antioxidant status in mice and rats. *J. Ethnopharmacol.* 97: 383-389.
- Loi MC, Poli F, Sacchetti G, Selenu MB, Ballero M (2004). Ethnopharmacology of Ogliastra (Villagrande Strisaili, Sardinia, Italy). *Fitoterapia* 75: 277-295.
- Malakov PY, Papanov GY, Boneva IM (1992). Neo-clerodane diterpenoids from *Teucrium montanum*. *Phytochem.* 31: 4029-4030.
- Malamas M, Marselos M (1992). The tradition of medicinal plants in Zagori, Epirus (Northwestern Greece). *J. Ethnopharmacol.* 37: 197-203.
- Merzouki A, Ed-derfoufi F, Molero Mesa J (2000). Contribution to the knowledge of Rifian traditional medicine. II: Folk medicine in Ksar Lakbir district (NW Morocco). *Fitoterapia* 71: 278-307.
- Miliauskas G, Venskutonis PR, van Beek TA (2004). Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food Chem.* 85: 231-237.
- Mulinacci N, Vincieri FF, Baldi A, Bambagiotti-Alberti M, Sendi A, Wagner H (1995). Flavonol glycosides from *Sedum telephium* subspecies *maximum* leaves. *Phytochem.* 38: 531-533.
- Muthu C, Ayyanar M, Raja N, Ignacimuthu S (2006). Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu, India. *J. Ethnobiol. Ethnomed.* 2: 43.
- Nahla AA (2003). Unique phenolic carboxylic acids from *Sanguisorba minor*. *Phytochem.* 63: 433-436.
- Nhu-Trang TT, Casabianca H, Grenier-Loustalot MF (2006). Deuterium/hydrogen ratio analysis of thymol, carvacrol, γ -terpinene and *p*-cymene in thyme, savory and oregano essential oils by gas chromatography-pyrolysis-isotope ratio mass spectrometry. *J. Chromatogr. A* 1132: 219-227.
- Nigro SA, Makunga NP, Grace OM (2004). Medicinal plants at the

- ethnobotany–biotechnology interface in Africa. *S. Afr. J. Bot.* 70: 89-96.
- Norscia I, Borgognini-Tarli SM (2006). Ethnobotanical reputation of plant species from two forests of Madagascar: A preliminary investigation. *S. Afr. J. Bot.* 72: 656-660.
- Novais MH, Santos I, Mendes S, Pinto-Gomes C (2004). Studies on pharmaceutical ethnobotany in Arrabida Natural Park (Portugal). *J. Ethnopharmacol.* 93: 183-195.
- Onanga M, Ekouya A, Ouabonzi A, Itoua GB (1999). Ethnobotanical, pharmacological and chemical studies of plants used in the treatment of 'Mwandza' dermatitis. *Fitoterapia* 70: 579-585.
- Otero R, Fonnegra R, Jimenez SL, Nunez V, Evans N, Alzate SP, Garca ME, Saldarriaga M, Del Valle G, Osorio RG, Daz A, Valderrama R, Duque A, Velez HN (2000). Snakebites and ethnobotany in the Northwest region of Colombia - Part I: Traditional use of plants. *J. Ethnopharmacol.* 71: 493-504.
- Oussalah M, Caillet S, Saucier L, Lacroix M (2006). Antimicrobial effects of selected plant essential oils on the growth of a *Pseudomonas putida* strain isolated from meat. *Meat Sci.* 73: 236-244.
- Oussalah M, Caillet S, Saucier L, Lacroix M (2007). Inhibitory effects of selected plant essential oils on the growth of four pathogenic bacteria: *E. coli* O157:H7, *Salmonella Typhimurium*, *Staphylococcus aureus* and *Listeria monocytogenes*. *Food Control* 18: 414-420.
- Palmese T, Rita M, Uncini Manganelli E, Tomei PE (2001). An ethnopharmacobotanical survey in the Sarrabus district (South-east Sardinia). *Fitoterapia* 72: 619-643.
- Passalacqua NG, Guarrera PM, De Fine G (2007). Contribution to the knowledge of the folk plant medicine in Calabria region (Southern Italy). *Fitoterapia* 78: 52-68.
- Perry EK, Pickering AT, Wang WW, J. Houghton, P, Perry NSL (1999). Medicinal plants and Alzheimer's disease: from ethnobotany to phytotherapy. *J. Pharm. Pharmacol.* 51: 527-534.
- PFaf (2006). Plants For A Future. Edible, medicinal and useful plants for a healthier world. (<http://www.pfaf.org/>; Accessed: December 23, 2006).
- Pieroni A, Nebel S, Quave C, Munz H, Heinrich M (2002). Ethnopharmacology of liakra: traditional weedy vegetables of the Arbereshe of the Vulture area in Southern Italy. *J. Ethnopharmacol.* 81: 165-185.
- Pieroni A, Quave C, Nebel S, Heinrich M (2002b). Ethnopharmacy of the ethnic Albanians (Arbëreshë) of Northern Basilicata, Italy. *Fitoterapia* 73: 217-241.
- Pieroni A, Giusti ME, Munz H, Lenzarini C, Turkovic G, Turkovic A (2003). Ethnobotanical knowledge of the Istro-Romanians of Zejane in Croatia. *Fitoterapia* 74: 710-719.
- Pieroni A, Quave CL (2005). Traditional pharmacopoeias and medicines among Albanians and Italians in Southern Italy: A comparison. *J. Ethnopharmacol.* 101: 258-270.
- Pieroni A, Muenz H, Akbulut M, Başer KHC, Durmuşkahya C (2005a). Traditional phytotherapy and trans-cultural pharmacy among Turkish migrants living in Cologne, Germany. *J. Ethnopharmacol.* 102: 69-88.
- Pieroni A, Dibra B, Grishaj G, Grishaj I, Maçai SG (2005b). Traditional phytotherapy of the Albanians of Lepushe, Northern Albanian Alps. *Fitoterapia* 76: 379-399.
- Rasoli I, Mirmostafa SA (2002). Antibacterial properties of *Thymus pubescens* and *Thymus serpyllum* essential oils. *Fitoterapia* 73: 244-250.
- Redzic S (1999). The syntaxonomical differentiation of the *Festuco-Brometea* Br.-Bl. and Tx. 1943 ex Klika and Hadač 1944 in the Balkans. *Ann. Bot.* 57: 167-180.
- Redzic S (2000). Patterns of succession of xerophylous vegetation on the Balkan. In "Vegetation science in retrospect and perspective" (eds White PS, Mucina L, Lepš J), Opulus press Uppsala pp. 76-79.
- Redzic SJ (2006a). Wild edible plants a their traditional use in human nutrition in Bosnia and Herzegovina. *Ecol. Food Nutr.* 45: 189-232.
- Redzic S (2006b). Organic wild production in Bosnia and Herzegovina. Proceedings of the 1st IFOAM International Conference on Organic Wild Production, Teslic, BiH. pp. 117-141.
- Redzic SS (2007). The ecological aspect of ethnobotany and ethnopharmacology of population in Bosnia and Herzegovina. *Coll. Antropol.* 31(3): 869-890.
- Redzic S (2009). Biodiversity of endemic plants as a source of new medicines (W. Balkan, SE Europe). *Planta Medica* 75(9): 988-988.
- Redzic S, Dalmatin M, Hamidovic M, Kadic J, Radevic M, Sevo Lj (2003). Biodiversity, geo-diversity and protection of natural and cultural heritages. National Environmental Action Plan of Bosnia and Herzegovina. Issue of Federation of BiH and RS, Sarajevo pp. 67-76.
- Redzic S, Lakusic R, Grujic-Vasic J, Tokic S, Kalinic D (1989). Medicinal plants in the ecosystems of the mountains Igman and Bjelasnica. *Lekovite sirovine/Medical, Drugs* 8: 5-14.
- Redzic S, Lakusic R, Tokic S (1991). Medicinal plants in some ecosystems of the Una river valley. *Bilten Društva ekologija BiH/Bulletin of Ecological Society of Bosnia and Herzegovina, Series B.* 6: 155-159.
- Redzic S, Grujic-Vasic J, Cerkez F, Mulabegovic N, Jaksic D, Ibrulj A (1997). Collection and utilization of medicinal plants during the war in Sarajevo. *Zeitschrift fur Phytotherapie*, Abstract band 4, Stuttgart pp. 23-24.
- Redzic S, Tuka M, Pajevic A (2006). Research into microscopic structure and essential oils of endemic medicinal plant species *Satureja subspicata* Bartl. ex Vis. (Lamiaceae). *Bosn J. Basic Med. Sci.* 6: 25-31.
- Rivera D, Obón C (1995). The ethnopharmacology of Madeira and Porto Santo Islands, a review. *J. Ethnopharmacol.* 46: 73-93.
- Sadikovic S (1929). *Liječenje biljem/Therapy by plants*. Publisher Svjetlost, (In Bosnian). Sarajevo pp. 5-424.
- Saikia AP, Ryakala VK, Sharma P, Goswami P, Bora U (2006). Ethnobotany of medicinal plants used by Assamese people for various skin ailments and cosmetics. *J. Ethnopharmacol.* 106: 149-157.
- Sajem AL, Gosai K (2006). Traditional use of medicinal plants by the Jaintia tribes in North Cachar Hills district of Assam, Northeast India. *J. Ethnobiol. Ethnomed.* 2: 33.
- Satou T, Mimaki Y, Kuroda M, Sashida Y, Hatakeyama Y (1996). A pyrrolone glucoside ester and steroidal saponins from *Lilium martagon*. *Phytochem.* 41: 1225-1230.
- Scherrer AM, Motti R, Weckerle CS (2005). Traditional plant use in the areas of Monte Vesole and Ascea, Cilento National Park (Campania, Southern Italy). *J. Ethnopharmacol.* 97: 129-143.
- Sheng-Ji P (2001). Ethnobotanical approaches of traditional medicine studies: some experiences from Asia. *Pharm. Biol.* 39(1): 74-79.
- Silic C (1984). Endemic plants. 1st edition. Publisher "Svjetlost" Sarajevo. (In Serbo-Croatian).
- Skočibušić M, Bezić N, Dunkić V (2006). Phytochemical composition and antimicrobial activities of the essential oils from *Satureja subspicata* Vis. growing in Croatia. *Food Chem.* 96: 20-28.
- Stobart A (2005). Medicinal Plants in Folk Tradition: An Ethnobotany of Britain and Ireland. *Soc. Hist. Med.* 18: 124-124.
- Testai L, Chericoni S, Calderone V, Nencioni G, Nieri P, Morelli I, Martinotti E (2002). Cardiovascular effects of *Urtica dioica* L. (Urticaceae) roots extracts: *in vitro* and *in vivo* pharmacological studies. *J. Ethnopharmacol.* 81: 105-109.
- Toker G, Küpeli E, Memisoğlu M, Yesilada E (2004). Flavonoids with antinociceptive and anti-inflammatory activities from the leaves of *Tilia argentea* (silver linden). *J. Ethnopharmacol.* 95: 393-397.
- Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Weeb DA (1980). *Flora Europea*. Cambridge University Press pp. 1-5.
- Uniyal SK, Singh KN, Jamwal P, Lal L (2006). Traditional use of medicinal plants among the tribal communities of Chhota Bhangal, Western Himalaya. *J. Ethnobiol. Ethnomed.* 2: 14.
- van-Wyk BEA (2002). Review of ethnobotanical research in Southern Africa. *S. Afr. J. Bot.* 68: 1-13.
- Vidyasagar GM, Prashantkumar P (2007). Traditional herbal remedies for gynecological disorders in women of Bidar district, Karnataka, India. *Fitoterapia* 78: 48-51.
- Vivienne W, Witkowski E, Balkwill K (2005). Application of diversity indices to appraise plant availability in the Traditional Medicinal markets of Johannesburg, S. Afri. *Biodiver. Conserv.* 14: 2971-3001.
- Werner L, Stevanovic-Janezic T, Spanoudaki M (1994). Cembratrienols and other components of white bark pine (*Pinus heldreichii*) oleoresin. *Phytochem.* 36: 1277-1279.